INDEX TO AUTHORS

AARONSON, STUART A. See STEPHENSON, TRONICK, REYNOLDS, and AARONSON, 427

AHMED, AFTAB, STRONG, DOUGLAS M., SELL, KENNETH W., THURMAN, GARY B., KNUDSEN, RICHARD C., WISTAR, RICHARD, JR., and GRACE, WILLIAM R. Demonstration of a blocking factor in the plasma and spinal fluid of patients with subacute sclerosing panencephalitis. I. Partial characterization, 902

ALLISON, A. C. See DAVIES, PAGE, and ALLISON, 1262

AMSBAUGH, DIANA F., HANSEN, CARL T., PRESCOTT, BENJAMIN, STASHAK, PHILIP W., ASOFSKY, RICHARD, and BAKER, PHILLIP J. Genetic control of the antibody response to Type III pneumococcal polysaccharide in mice. II. Relationship between IgM immunoglobulin levels and the ability to give an IgM antibody response, 1499

ANDERSON, CLARK L., and GREY, HOWARD M. Receptors for aggregated IgG on mouse lymphocytes. Their presence on thymocytes, thymus-derived, and bone marrow-derived lymphocytes, 1175

ANDERSON, SETH E., JR., and REMINGTON, JACK S. Effect of normal and activated human macrophages on *Toxoplasma gondii*, 1154

ARMERDING, DIETER, and KATZ, DAVID H. Activation of T and B lymphocytes in vitro. I. Regulatory influence of bacterial lipopolysaccharide (LPS) on specific T-cell helper function, 24

ASOFSKY, RICHARD. See AMSBAUGH, HANSEN, PRESCOTT, STASHAK, ASOFSKY, and BAKER, 1499

—. See MORSE, HARRISON, and ASOFSKY, 721

AULT, KENNETH A., and UNANUE, EMIL R. Events after the binding of antigen to lymphocytes. Removal and regeneration of the antigen receptor, 1110

—. See UNANUE, AULT, and KARNOVSKY, 295

AUSTEN, K. FRANK. See FEARON, AUSTEN, and RUDDY, 355

BAKER, PHILLIP J. See AMSBAUGH, HANSEN, PRESCOTT, STASHAK, ASOFSKY, AND BAKER, 1499

BARONOWSKY, P. E. See DEVLIN, MCCURDY, and BARONOWSKY, 224, 230

BEHIN, REZA. See DOYLE, BEHIN, MAUEL, and ROWE, 1061

BENACERRAF, BARUJ. See KATZ, HAMAOKA, and BENACERRAF, 1464

BENNETT, MICHAEL. See KUMAR, BENNETT, and ECKNER, 1093

BERKO, BARBARA. See SHAW, BERKO, and WEGMANN, 767

BEVAN, MICHAEL J., EPSTEIN, RUTH, and COHN, MELVIN. The effect of 2-mercaptoethanol on murine mixed lymphocyte cultures, 1025

BIGLEY, ROBERT H., and STANKOVA, LIBUSE. Uptake and reduction of oxidized and reduced ascorbate by human leukocytes, 1084

BILLINGS, PAUL. See KAGNOFF, BILLINGS, and COHN, 407

BINZ, HANS, LINDENMANN, JEAN, and WIGZELL, HANS. Cell-bound receptors for alloantigens on normal lymphocytes. I. Characterization of receptor-carrying cells by the use of antibodies to alloantibodies, 877

BISMUTH, A. See NEAUPORT-SAUTES, BISMUTH, KOURILSKY, and MANUEL, 957

BITTER-SUERMANN, DIETER. See DUKOR, SCHUMANN, GISLER, DIERICH, KÖNIG, HADDING, and BITTER-SUERMANN, 337
BOKISCH, VIKTOR A. See THEOFILOPOULOS, BOKISCH, and DIXON, 696
BOSMA, GAYLE C. See BOSMA and BOSMA, 512
BOSMA, MELVIN J., and BOSMA, GAYLE C. Congenic mouse strains: the expression of a hidden immunoglobulin allotype in a congenic partner strain of BALB/c mice, 512
BRAUN, DIETMAR G. See CRAMER and BRAUN, 1513
BRO-JØRGENSEN, KLAUS. See VOLKERT, MARKER, and BRO-JØRGENSEN, 1329
BROWN, DAVID M. See LEE, MAUER, BROWN, SUTHERLAND, MICHAEL, and NAJARIAN, 793
BULLOCK, WESLEY W. See COUTINHO, GRONOWICZ, BULLOCK, and MÖLLER, 74
BUTLER, VINCENT P., JR. See SCHMIDT, KAUFMAN, and BUTLER, 278

CAMPBELL, STEPHEN. See KAGNOFF and CAMPBELL, 398
CAPPEL, ROGER, SCHLUEDERBERG, ANN, GIFFORD, ROBERT H., and HORSTMANN, DOROTHY M. Characterization of a human acute phase protein found in association with rubella virus infection, 497
CAPRA, J. D. See KUNKEL, WINCHESTER, JOSLIN, and CAPRA, 128
CARSON, DENNIS. See WEIGERT, RASCHKE, CARSON, and COHN, 137
CARTER, PHILIP B., and COLLINS, FRANK M. The route of enteric infection in normal mice, 1189
CEBRA, JOHN J. See JONES, CEBRA, and HERZENBERG, 581
CHOI, TAT K., SLEIGHT, DOUGLAS R., and NISONOFF, ALFRED. General method for isolation and recovery of B cells bearing specific receptors, 761
CHUSED, T. M. See KIRCHNER, CHUSED, HERBERMAN, HOLDEN, and LAVRIN, 1473
CISAR, JOHN, KABAT, ELVIN A., LIAO, JERRY, and POTTER, MICHAEL. Immunochemical studies on mouse myeloma proteins reactive with dextrans or with fructosans and on human antilevans, 159
CLAFLIN, J. LATHAM, LIEBERMAN, ROSE, and DAVIE, JOSEPH M. Clonal nature of the immune response to phosphorylcholine. I. Specificity, class, and idiotype of phosphorylcholine-binding receptors on lymphoid cells, 58
CLAGETT, JAMES A., and WEIGLE, WILLIAM O. Roles of T and B lymphocytes in the termination of unresponsiveness to autologous thyroglobulin in mice, 643
CLICK, ROBERT E. Immune responses in vitro. IX. Absence of the MLR-stimulatory, M-locus product from thymus cells, 1628
COHN, MELVIN. See BEVAN, EPSTEIN, and COHN, 1025
—. See KAGNOFF, BILLINGS, and COHN, 407
—. See WEIGERT, RASCHKE, CARSON, and COHN, 137
COHN, ZANVIL A. See GORDON, TODD, and COHN, 1228
—. See STEINMAN and COHN, 380
—. See STEINMAN, LUSTIG, and COHN, 1431
COLLINS, FRANK M. See CARTER and COLLINS, 1189
—. See VOLKMAN and COLLINS, 264
COONS, ALBERT H. See RUBIN, MACDONALD, and COONS, 439
COUTINHO, ANTONIO, GRONOWICZ, EVA, BULLOCK, WESLEY W., and MÖLLER, GÖRAN. Mechanism of thymus-independent immunocyte triggering. Mitogenic activation of

B cells results in specific immune responses, 74
CRAIGHEAD, JOHN E., and HIGGINS, DUNCAN A. Genetic influences affecting the occurrence of a diabetes mellitus-like disease in mice infected with the encephalomyocarditis virus, 414
CRAMER, MATTHIAS, and BRAUN, DIETMAR G. Genetics of restricted antibodies to streptococcal group polysaccharides in mice. I. Strain differences of isoelectric focusing spectra of group A hyperimmune antisera, 1513
CROCE, CARLO M., and KOPROWSKI, HILARY. Concordant segregation of the expression of SV40 T antigen and human chromosome 7 in mouse-human hybrid subclones, 1350

DAHL, LEWIS K., and TUTHILL, RALPH. Further evidence of the toxicity of NaCl. Increased blood pressure and mortality in the spontaneously hypertensive rat, 617
DAVIE, JOSEPH M. See CLAFLIN, LIEBERMAN, and DAVIE, 58
DAVIES, PHILIP, PAGE, ROY C., and ALLISON, A. C. Changes in cellular enzyme levels and extracellular release of lysosomal acid hydrolases in macrophages exposed to group A streptococcal cell wall substance, 1262
DECARY, FRANCINE. See RUBINSTEIN, LIU, STREUN, and DECARY, 313
DEVLIN, R. G., MCCURDY, J. D., and BARONOWSKY, P. E. Mixed lymphocyte reactivity against normal cells by splenic lymphocytes from tumor-bearing mice. I. Studies of vigorous immune responsiveness induced in F_1 mice by parental strain tumor cells, 224
—, —, and —. II. Studies of autoimmune-like activity in completely syngeneic and semisyngeneic systems, 230
DIERICH, MANFRED. See DUKOR, SCHUMANN, GISLER, DIERICH, KÖNIG, HADDING, and BITTER-SUERMANN, 337
DINARELLO, CHARLES A., GOLDIN, NATHAN P., and WOLFF, SHELDON M. Demonstration and characterization of two distinct human leukocytic pyrogens, 1369
DI PAULI, RAIMUND. See KOLB, DI PAULI, and WEILER, 467
DIXON, FRANK J. See THEOFILOPOULOS, BOKISCH, and DIXON, 696
DORSETT, BRENT H. See IOACHIM, KELLER, DORSETT, and PEAR[illegible], 1382
DOYLE, JOHN J., BEHIN, REZA, MAUEL, JACQUES, and ROWE, DAVID S. Antibody-induced movement of membrane components of *Leishmania enrietii*, 1061
DUKOR, PETER, SCHUMANN, GEBHARD, GISLER, ROLAND H., DIERICH, MANFRED, KÖNIG, WOLFGANG, HADDING, ULRICH, and BITTER-SUERMANN, DIETER. Complement-dependent B-cell activation by cobra venom factor and other mitogens? 337

ECKNER, ROBERT J. See KUMAR, BENNETT, and ECKNER, 1093
EDELMAN, GERALD M. See SPEAR and EDELMAN, 249
ELFENBEIN, GERALD J. See GELFAND, ELFENBEIN, FRANK, and PAUL, 1125
EPSTEIN, RUTH. See BEVAN, EPSTEIN, and COHN, 1025
ESSELMAN, W. J., and MILLER, H. C. Brain and thymus lipid inhibition of antibrain-associated θ-cytotoxicity, 445

FARIS, BARBARA. See ROWE, MCGOODWIN, MARTIN, SUSSMAN, GRAHN, FARIS, and FRANZBLAU, 180
FEARON, DOUGLAS T., AUSTEN, K. FRANK, and RUDDY, SHAUN. Pro-

perdin factor D: characterization of its active site and isolation of the precursor form, 355

FLEISSNER, ERWIN. See YOSHIKI, MELLORS, HARDY, and FLEISSNER, 925

FORRISTAL, JUDITH. See VALLOTA, GÖTZE, SPIEGELBERG, FORRISTAL, WEST, and MÜLLER-EBERHARD, 1249

FRANK, MICHAEL M. See GELFAND, ELFENBEIN, FRANK, and PAUL, 1125

FRANZBLAU, CARL. See ROWE, MCGOODWIN, MARTIN, SUSSMAN, GRAHN, FARIS, and FRANZBLAU, 180

FRENKEL, J. K. See HOFF and FRENKEL, 560

FU, SHU MAN, WINCHESTER, ROBERT J., and KUNKEL, HENRY G. Occurrence of surface IgM, IgD, and free light chains on human lymphocytes, 451

FUCHS, SARA, MOZES, EDNA, MAOZ, ARIEH, and SELA, MICHAEL. Thymus independence of a collagen-like synthetic polypeptide and of collagen, and the need for thymus and bone marrow cell cooperation in the immune response to gelatin, 148

GALARDY, R. E. See KRAEHENBUHL, GALARDY, and JAMIESON, 208

GARVIN, A. J., SPICER, S. S., PARMLEY, R. T., and MUNSTER, A. M. Immunohistochemical demonstration of IgG in Reed-Sternberg and other cells in Hodgkin's disease, 1077

GELFAND, MICHAEL C., ELFENBEIN, GERALD J., FRANK, MICHAEL M., and PAUL, WILLIAM E. Ontogeny of B lymphocytes. II. Relative rates of appearance of lymphocytes bearing surface immunoglobulin and complement receptors, 1125

—, SACHS, DAVID H., LIEBERMAN, ROSE, and PAUL, WILLIAM E. Ontogeny of B lymphocytes. III. *H-2* linkage of a gene controlling the rate of appearance of complement receptor lymphocytes, 1142

GEORGE, W. J. See IGNARRO, LINT, and GEORGE, 1395

GERBASE-DELIMA, MARIA. See TITTOR, GERBASE-DELIMA, and WALFORD, 1488

GIFFORD, ROBERT H. See CAPPEL, SCHLUEDERBERG, GIFFORD, and HORSTMANN, 497

GISLER, ROLAND H. See DUKOR, SCHUMANN, GISLER, DIERICH, KÖNIG, HADDING, and BITTER-SUERMANN, 337

GLOVER, ROBYN G. See OATES, STOKES, STOREY, GLOVER, and SNOW, 239

GOLDIN, NATHAN P. See DINARELLO, GOLDIN, and WOLFF, 1369

GÖTZE, OTTO, and MÜLLER-EBERHARD, HANS J. The role of properdin in the alternate pathway of complement activation, 44

—. See VALLOTA, GÖTZE, SPIEGELBERG, FORRISTAL, WEST, and MÜLLER-EBERHARD, 1249

GOLUB, EDWARD S. See VAN DEN ENGH and GOLUB, 1621

GORDON, SAIMON, TODD, JEAN, and COHN, ZANVIL A. In vitro synthesis and secretion of lysozyme by mononuclear phagocytes, 1228

—. See UNKELESS, GORDON, and REICH, 834

GRACE, WILLIAM R. See AHMED, STRONG, SELL, THURMAN, KNUDSEN, WISTAR, and GRACE, 902

GRAHN, DOUGLAS. See ROWE, MCGOODWIN, MARTIN, SUSSMAN, GRAHN, FARIS, and FRANZBLAU, 180

GRAW, CAROLE M. See LOHRMANN, GRAW, and GRAW, 1037

GRAW, ROBERT G., JR. See LOHRMANN, GRAW, and GRAW, 1037

—. See LOHRMANN, NOVIKOVS, and GRAW, 1553

GREEN, IRA. See SHEVACH, GREEN, and PAUL, 679
—. See SHEVACH, PAUL, and GREEN, 661
GREY, HOWARD M. See ANDERSON and GREY, 1175
GRIFFIN, FRANK M., JR., and SILVERSTEIN, SAMUEL C. Segmental response of the macrophage plasma membrane to a phagocytic stimulus, 323
GRONOWICZ, EVA. See COUTINHO, GRONOWICZ, BULLOCK, and MÖLLER, 74
GRUNDKE-IQBAL, INGE. See VITETTA, GRUNDKE-IQBAL, HOLMES, and UHR, 862

HA, TAI-YOU, WAKSMAN, BYRON H., and TREFFERS, H. PETER. The thymic suppressor cell. I. Separation of subpopulations with suppressor activity, 13
HADDING, ULRICH. See DUKOR, SCHUMANN, GISLER, DIERICH, KÖNIG, HADDING, and BITTER-SUERMANN, 337
HAMAOKA, TOSHIYUKI, and KATZ, DAVID H. Immunological tolerance in bone marrow-derived lymphocytes. II. Effects of allogeneic cell interactions and enzymatic digestion with trypsin on inactivated hapten-specific precursors of antibody-forming cells, 1446
—. See KATZ, HAMAOKA, and BENACERRAF, 1464
HANNA, M. G., JR. See IHLE, HANNA, ROBERSON, and KENNEY, 1568
HANSEN, CARL T. See AMSBAUGH, HANSEN, PRESCOTT, STASHAK, ASOFSKY, and BAKER, 1499
HARDY, WILLIAM D., JR. See YOSHIKI, MELLORS, HARDY, and FLEISSNER, 925
HARRIS, ALAN W. See STOCKER, MARCHALONIS, and HARRIS, 785
HARRISON, MICHAEL R. See MORSE, HARRISON, and ASOFSKY, 721
HERBERMAN, R. B. See KIRCHNER, CHUSED, HERBERMAN, HOLDEN, and LAVRIN, 1473
HERZENBERG, LEONARD A. See JONES, CEBRA, and HERZENBERG, 581
HIGGINS, DUNCAN A. See CRAIGHEAD and HIGGINS, 414
HOFF, RICHARD L., and FRENKEL, J. K. Cell-mediated immunity against *Besnoitia* and *Toxoplasma* in specifically and cross-immunized hamsters and in cultures, 560
HOLDEN, H. T. See KIRCHNER, CHUSED, HERBERMAN, HOLDEN, and LAVRIN, 1473
HOLMES, KATHRYN V. See VITETTA, GRUNDKE-IQBAL, HOLMES, and UHR, 862
HOLMGREN, J., LINDHOLM, L., and LÖNNROTH, I. Interaction of cholera toxin and toxin derivatives with lymphocytes. I. Binding properties and interference with lectin-induced cellular stimulation, 801
HORSTMANN, DOROTHY M. See CAPPEL, SCHLUEDERBERG, GIFFORD, and HORSTMANN, 497
HÜNIG, T., SCHIMPL, A., and WECKER, E. Autoradiographic studies on the proliferation of antibody-producing cells in vitro, 754
HUMPHREY, W., JR. See LIEBERMAN, POTTER, MUSHINSKI, HUMPHREY, and RUDIKOFF, 983
HUNT, SIMON V., and WILLIAMS, ALAN F. The origin of cell surface immunoglobulin of marrow-derived and thymus-derived lymphocytes of the rat, 479
HUSBY, G., NATVIG, J. B., and SLETTEN, K. New, third class of amyloid fibril protein, 773

IGNARRO, L. J., LINT, T. F., and GEORGE, W. J. Hormonal control of lysosomal enzyme release from human neutrophils. Effects of autonomic agents on enzyme release, phagocytosis, and cyclic nucleotide levels, 1395
IHLE, J. N., HANNA, M. G., JR.,

ROBERSON, L. E., and KENNEY, F. T. Autogenous immunity to endogenous RNA tumor virus. Identification of antibody reactivity to select viral antigens, 1568

IOACHIM, HARRY L., KELLER, STEVEN E., DORSETT, BRENT H., and PEARSE, ANTONIA. Induction of partial immunologic tolerance in rats and progressive loss of cellular antigenicity in gross virus lymphoma, 1382

ISERSKY, CHAVIVA. See KULCZYCKI, ISERSKY, and METZGER, 600

ISHIBASHI, T. See MACKANESS, LAGRANGE, and ISHIBASHI, 1540

—. See MACKANESS, LAGRANGE, MILLER, and ISHIBASHI, 543

JAMIESON, J. D. See KRAEHENBUHL, GALARDY, and JAMIESON, 208

JOHNSON, B. M. See MOSIER, JOHNSON, PAUL, and MCMASTER, 1354

JONES, PATRICIA P., CEBRA, JOHN J., and HERZENBERG, LEONARD A. Restriction of gene expression in B lymphocytes and their progeny. I. Commitment to immunoglobulin allotype, 581

JOSLIN, F. G. See KUNKEL, WINCHESTER, JOSLIN, and CAPRA, 128

KABAT, ELVIN A. See CISAR, KABAT, LIAO, and POTTER, 159

KAGNOFF, MARTIN F., BILLINGS, PAUL, and COHN, MELVIN. Functional characteristics of Peyer's patch lymphoid cells. II. Lipopolysaccharide is thymus dependent, 407

— and CAMPBELL, STEPHEN. Functional characteristics of Peyer's patch lymphoid cells. I. Induction of humoral antibody and cell-mediated allograft reactions, 398

KAPLAN, JOSEPH, SHOPE, THOMAS C., and PETERSON, WARD D., JR. Epstein-Barr virus-negative human malignant T-cell lines, 1070

KARLSSON, M. See O'TOOLE, STEJSKAL, PERLMANN, and KARLSSON, 457

KARNOVSKY, MORRIS J. See SCHNEEBERGER, LEBER, KARNOVSKY, and MCCLUSKEY 1283

—. See UNANUE, AULT, and KARNOVSKY, 295

KATZ, DAVID H., HAMAOKA, TOSHIYUKI, and BENACERRAF, BARUJ. Immunological tolerance in bone marrow-derived lymphocytes. III. Tolerance induction in primed B cells by hapten conjugates of unrelated immunogenic or "nonimmunogenic" carriers, 1464

—. See ARMERDING and KATZ, 24

—. See HAMAOKA and KATZ, 1446

KAUFMAN, BETTE M. See SCHMIDT, KAUFMAN, and BUTLER, 278

KELLER, STEVEN E. See IOACHIM, KELLER, DORSETT, and PEARSE, 1382

KENNEL, STEPHEN J. The K-chains of the immunoglobulin from a continuous culture of human lymphocytes (Daudi) have an unusual molecular size, 1031

KENNEY, F. T. See IHLE, HANNA, ROBERSON, and KENNEY, 1568

KIMURA, ARTHUR K. Inhibition of specific cell-mediated cytotoxicity by anti-T-cell receptor antibody, 888

KINDRED, BERENICE, and LOOR, FRANCIS. Activity of host-derived T cells which differentiate in nude mice grafted with co-isogenic or allogeneic thymuses, 1215

KIRCHNER, H., CHUSED, T. M., HERBERMAN, R. B., HOLDEN, H. T., and LAVRIN, D. H. Evidence of suppressor cell activity in spleens of mice bearing primary tumors induced by Moloney sarcoma virus, 1473

KNUDSEN, RICHARD C. See AHMED,

STRONG, SELL, THURMAN, KNUDSEN, WISTAR, and GRACE, 902

KÖNIG, WOLFGANG. See DUKOR, SCHUMANN, GISLER, DIERICH, KÖNIG, HADDING, and BITTER-SUERMANN, 337

KOLB, CORNELIA, DI PAULI, RAIMUND, and WEILER, EBERHARDT. Induction of IgG by lipid A in the newborn mouse, 467

KOOK, ABRAHAM I., and TRAININ, NATHAN. Hormone-like activity of a thymus humoral factor on the induction of immune competence in lymphoid cells, 193

KOPROWSKI, HILARY. See CROCE and KOPROWSKI, 1350

KOURILSKY, F. M. See NEAUPORT-SAUTES, BISMUTH, KOURILSKY, and MANUEL, 957

KRAEHENBUHL, J. P., GALARDY, R. E., and JAMIESON, J. D. Preparation and characterization of an immunoelectron microscope tracer consisting of a heme-octapeptide coupled to Fab, 208

KULCZYCKI, ANTHONY, JR., ISERSKY, CHAVIVA, and METZGER, HENRY. The interaction of IgE with rat basophilic leukemia cells. I. Evidence for specific binding of IgE, 600

KUMAR, VINAY, BENNETT, MICHAEL, and ECKNER, ROBERT J. Mechanisms of genetic resistance to Friend virus leukemia in mice. I. Role of ^{89}Sr-sensitive effector cells responsible for rejection of bone marrow allografts, 1093

KUNKEL, HENRY G., WINCHESTER, ROBERT J., JOSLIN, F. G., and CAPRA, J. D. Similarities in the light chains of anti-γ-globulins showing cross-idiotypic specificities, 128

—. See FU, WINCHESTER, and KUNKEL, 451

—. See UTERMOHLEN, WINFIELD, ZABRISKIE, and KUNKEL, 1019

LAGRANGE, P. H., MACKANESS, G. B., and MILLER, T. E. Influence of dose and route of antigen injection on the immunological induction of T cells, 528

—, —, and —. Potentiation of T-cell-mediated immunity by selective suppression of antibody formation with cyclophosphamide, 1529

—. See MACKANESS, LAGRANGE, and ISHIBASHI, 1540

—. See MACKANESS, LAGRANGE, MILLER and ISHIBASHI, 543

LAMM, MICHAEL E. See POGER and LAMM, 629

LAVRIN, D. H. See KIRCHNER, CHUSED, HERBERMAN, HOLDEN, and LAVRIN, 1473

LAWRENCE, DAVID A., and WEIGLE, WILLIAM O. Stimulation of antibody production to the hapten 2,4-dinitrobenzene by affinity labeled murine lymphoid cells. I. Ability of affinity labeled murine lymphoid cells to activate the in vivo immune response, 943

LEBER, P. D. See SCHNEEBERGER, LEBER, KARNOVSKY, and MCCLUSKEY, 1283

LEE, CHUE SHUE, MAUER, S. MICHAEL, BROWN, DAVID M., SUTHERLAND, DAVID E. R., MICHAEL, ALFRED F., and NAJARIAN, JOHN S. Renal transplantation in diabetes mellitus in rats, 793

LIAO, JERRY. See CISAR, KABAT, LIAO, and POTTER, 159

LIEBERMAN, ROSE, POTTER, MICHAEL, MUSHINSKI, E. B., HUMPHREY, W., JR., and RUDIKOFF, S. Genetics of a new IgV_H(T15 idiotype)marker in the mouse regulating natural antibody to phosphorylcholine, 983

—. See CLAFLIN, LIEBERMAN, and DAVIE, 58

—. See GELFAND, SACHS, LIEBERMAN, and PAUL, 1142

LIEW, F. Y., and PARISH, C. R. Lack of a correlation between cell-mediated immunity to the carrier

and the carrier-hapten helper effect, 779
LINDENMANN, JEAN. See BINZ, LINDENMANN, and WIGZELL, 877
LINDHOLM, L. See HOLMGREN, LINDHOLM, and LÖNNROTH, 801
LINT, T. F. See IGNARRO, LINT, and GEORGE, 1395
LIU, NGUKOY. See RUBINSTEIN, LIU, STREUN, and DECARY, 313
LOEB, JOHN N. See RIFKIN, LOEB, MOORE, and REICH, 1317
LÖNNROTH, I. See HOLMGREN, LINDHOLM, and LÖNNROTH, 801
LOGIE, PAMELA S. See MCGREGOR and LOGIE, 1415
LOHRMANN, HANS-PETER, GRAW, CAROLE M., and GRAW, ROBERT G., JR. Stimulated lymphocyte cultures. Responder recruitment and cell cycle kinetics, 1037
—, NOVIKOVS, LIGITA, and GRAW, ROBERT G., JR. Cellular interactions in the proliferative response of human T and B lymphocytes to phytomitogens and allogeneic lymphocytes, 1553
LOOR, FRANCIS. See KINDRED and LOOR, 1215
LOZNER, EUGENE C., SACHS, DAVID H., and SHEARER, GENE M. Genetic control of the immune response to staphylococcal nuclease. I. *Ir*-Nase: control of the antibody response to nuclease by the *Ir* region of the mouse *H-2* complex, 1204
LUSTIG, DINAH S. See STEINMAN, LUSTIG, and COHN, 1431

MACDONALD, A. BRUCE. See RUBIN, MACDONALD, and COONS, 439
MACKANESS, G. B., LAGRANGE, P. H., and ISHIBASHI, T. The modifying effect of BCG on the immunological induction of T cells, 1540
—, —, MILLER, T. E., and ISHIBASHI, T. Feedback inhibition of specifically sensitized lymphocytes, 543
MACKANESS, G. B. See LAGRANGE, MACKANESS, and MILLER, 528, 1529
MANUEL, Y. See NEAUPORT-SAUTES, BISMUTH, KOURILSKY, and MANUEL, 957
MAOZ, ARIEH. See FUCHS, MOZES, MAOZ, and SELA, 148
MARCHALONIS, JOHN J. See STOCKER, MARCHALONIS, and HARRIS, 785
MARKER, OLE. See VOLKERT, MARKER, and BRO-JØRGENSEN, 1329
MARTIN, GEORGE R. See ROWE, MCGOODWIN, MARTIN, SUSSMAN, GRAHN, FARIS, and FRANZBLAU, 180
MAUEL, JACQUES. See DOYLE, BEHIN, MAUEL, and ROWE, 1061
MAUER, S. MICHAEL. See LEE, MAUER, BROWN, SUTHERLAND, MICHAEL, and NAJARIAN, 793
MCCLUSKEY, R. T. See SCHNEEBERGER, LEBER, KARNOVSKY, and MCCLUSKEY, 1283
MCCURDY, J. D. See DEVLIN, MCCURDY, and BARONOWSKY, 224, 230
MCGOODWIN, ERMONA B. See ROWE, MCGOODWIN, MARTIN, SUSSMAN, GRAHN, FARIS, and FRANZBLAU, 180
MCGREGOR, D. D., and LOGIE, PAMELA S. The mediator of cellular immunity. VII. Localization of sensitized lymphocytes in inflammatory exudates, 1415
MCMASTER, P. R. B. See MOSIER, JOHNSON, PAUL, and MCMASTER, 1354
MCMICHAEL, A. J., and WILLIAMSON, A. R. Clonal memory. I. Time-course of proliferation of B-memory cells, 1361
MELLORS, ROBERT C. See YOSHIKI, MELLORS, HARDY, and FLEISSNER, 925
MELSOM, HAAKON. Cytotoxic activity of mouse macrophages by various inhibitors, 1049

METZGER, HENRY. See KULCZYCKI, ISERSKY, and METZGER, 600
MICHAEL, ALFRED F. See LEE, MAUER, BROWN, SUTHERLAND, MICHAEL, and NAJARIAN, 793
MILLER, H. C. See ESSELMAN and MILLER, 445
MILLER, J. F. A. P. See SPRENT and MILLER, 1
MILLER, T. E. See LAGRANGE, MACKANESS, and MILLER, 528, 1529
—. See MACKANESS, LAGRANGE, MILLER, and ISHIBASHI, 543
MÖLLER, GÖRAN. Effect of B-cell mitogens on lymphocyte subpopulations possessing C'3 and Fc receptors, 969
—. See COUTINHO, GRONOWICZ, BULLOCK, and MÖLLER, 74
MOORE, GEORGE. See RIFKIN, LOEB, MOORE, and REICH, 1317
MORSE, HERBERT C., III, HARRISON, MICHAEL R., and ASOFSKY, RICHARD. Graft-vs.-host reactions in reciprocal hybrid mice. I. Dissociation of T-cell activities in the mixed lymphocyte reaction and two graft-vs.-host assays, 721
MOSIER, D. E., JOHNSON, B. M., PAUL, WILLIAM E., and MCMASTER, P. R. B. Cellular requirements for the primary in vitro antibody response to DNP-Ficoll, 1354
MOZES, EDNA. See FUCHS, MOZES, MAOZ, and SELA, 148
MÜLLER-EBERHARD, HANS J. See GÖTZE and MÜLLER-EBERHARD, 44
—. See VALLOTA, GÖTZE, SPIEGELBERG, FORRISTAL, WEST, and MÜLLER-EBERHARD, 1249
MUNSTER, A. M. See GARVIN, SPICER, PARMLEY, and MUNSTER, 1077
MUSHINSKI, E. B. See LIEBERMAN, POTTER, MUSHINSKI, HUMPHREY, and RUDIKOFF, 983

NAJARIAN, JOHN S. See LEE, MAUER, BROWN, SUTHERLAND, MICHAEL, and NAJARIAN, 793
NATVIG, J. B. See HUSBY, NATVIG, and SLETTEN, 773
NEAUPORT-SAUTES, C., BISMUTH, A., KOURILSKY, F. M., and MANUEL, Y. Relationship between HL-A antigens and β_2-microglobulin as studied by immunofluorescence on the lymphocyte membrane, 957
NISONOFF, ALFRED. See CHOI, SLEIGHT, and NISONOFF, 761
NOSSAL, G. J. V. See SCHRADER and NOSSAL, 1582
NOVIKOVS, LIGITA. See LOHRMANN, NOVIKOVS, and GRAW, 1553

OATES, HELEN F., STOKES, GORDON S., STOREY, BRIAN G., GLOVER, ROBYN G., and SNOW, BEVERLEY F. Renal hypertension in rats immunized against angiotensin I and angiotensin II, 239
O'TOOLE, C., STEJSKAL, V., PERLMANN, P., and KARLSSON, M. Lymphoid cells mediating tumor-specific cytotoxicity to carcinoma of the urinary bladder. Separation of the effector population using a surface marker, 457

PAGE, ROY C. See DAVIES, PAGE, and ALLISON, 1262
PARISH, C. R. See LIEW and PARISH, 779
PARKER, CHARLES W. See SHEARER, PHILPOTT, and PARKER, 367
PARMLEY, R. T. See GARVIN, SPICER, PARMLEY, and MUNSTER, 1077
PAUL, WILLIAM E. See GELFAND, ELFENBEIN, FRANK, and PAUL, 1125
—. See GELFAND, SACHS, LIEBERMAN, and PAUL, 1142
—. See MOSIER, JOHNSON, PAUL, and MCMASTER, 1354
—. See SHEVACH, GREEN, and PAUL, 679

PAUL, WILLIAM E. See SHEVACH, PAUL, and GREEN, 661
PEARSE, ANTONIA. See IOACHIM, KELLER, DORSETT, and PEARSE, 1382
PERLMANN, P. See O'TOOLE, STEJSKAL, PERLMANN, and KARLSSON, 457
PETERSON, WARD D., JR. See KAPLAN, SHOPE, and PETERSON, 1070
PHILPOTT, GORDON W. See SHEARER, PHILPOTT, and PARKER, 367
PLATE, JANET M. D. Mixed lymphocyte culture responses of mice. Genetic analysis of the responses to *H-2D*^{d} specificities, 851
POGER, MARSHALL E., and LAMM, MICHAEL E. Localization of free and bound secretory component in human intestinal epithelial cells. A model for the assembly of secretory IgA, 629
POTTER, MICHAEL. See CISAR, KABAT, LIAO, and POTTER, 159
—. See LIEBERMAN, POTTER, MUSHINSKI, HUMPHREY, and RUDIKOFF, 983
PRENDERGAST, ROBERT A. See WILLENBORG and PRENDERGAST, 820
PRESCOTT, BENJAMIN. See AMSBAUGH, HANSEN, PRESCOTT, STASHAK, ASOFSKY, and BAKER, 1499
PROSS, HUGH. See SCHIRRMACHER, RUBIN, PROSS, and WIGZELL, 93

RASCHKE, WILLIAM C. See WEIGERT, RASCHKE, CARSON, and COHN, 137
REICH, E. See RIFKIN, LOEB, MOORE, and REICH, 1317
—. See UNKELESS, GORDON, and REICH, 834
REMINGTON, JACK S. See ANDERSON and REMINGTON, 1154
REYNOLDS, ROBERTA K. See STEPHENSON, TRONICK, REYNOLDS, and AARONSON, 427
RIFKIN, DANIEL B., LOEB, JOHN N., MOORE, GEORGE, and REICH, E. Properties of plasminogen activators formed by neoplastic human cell cultures, 1317
ROBERSON, L. E. See IHLE, HANNA, ROBERSON, and KENNEY, 1568
RODKEY, L. SCOTT. Studies of idiotypic antibodies. Production and characterization of autoantiidiotypic antisera, 712
ROSENSTREICH, DAVID L. See VAN BOXEL and ROSENSTREICH, 1002
ROWE, DAVID S. See DOYLE, BEHIN, MAUEL, and ROWE, 1061
ROWE, DAVID W., MCGOODWIN, ERMONA B., MARTIN, GEORGE R., SUSSMAN, MICHAEL D., GRAHN, DOUGLAS, FARIS, BARBARA, and FRANZBLAU, CARL. A sex-linked defect in the cross-linking of collagen and elastin associated with the mottled locus in mice, 180
RUBIN, ARNOLD S., MACDONALD, A. BRUCE, and COONS, ALBERT H. Specific heterologous enhancement of immune responses. VI. Partial purification of a nonspecific enhancing factor from supernates of allogeneically stimulated human lymphocyte cell lines, 439
RUBIN, BENT, and WIGZELL, HANS. Lymphocyte proliferation in vitro induced by hapten autologous protein conjugates. I. A study on the class of lymphocytes responding in vitro and on the nature and specificity of their receptors, 732
—. See SCHIRRMACHER, RUBIN, PROSS, and WIGZELL, 93
RUBINSTEIN, PABLO, LIU, NGUKOY, STREUN, EDWIN W., and DECARY, FRANCINE. Electrophoretic mobility and agglutinability of red blood cells: a "new" polymorphism in mice, 313
RUDDY, SHAUN. See FEARON, AUSTEN, and RUDDY, 355
RUDIKOFF, S. See LIEBERMAN, POTTER, MUSHINSKI, HUMPHREY, and RUDIKOFF, 983

SACHS, DAVID H. See GELFAND, SACHS, LIEBERMAN, and PAUL, 1142
—. See LOZNER, SACHS, and SHEARER, 1204
SCHIMPL, A. See HÜNIG, SCHIMPL, and WECKER, 754
SCHIRRMACHER, VOLKER, RUBIN, BENT, PROSS, HUGH, and WIGZELL, HANS. Cytotoxic immune cells with specificity for defined soluble antigens. IV. Antibody as mediator of specific cytotoxicity, 93
SCHLUEDERBERG, ANN. See CAPPEL, SCHLUEDERBERG, GIFFORD, and HORSTMANN, 497
SCHMIDT, DONALD H., KAUFMAN, BETTE M., and BUTLER, VINCENT P., JR. Persistence of hapten-antibody complexes in the circulation of immunized animals after a single intravenous injection of hapten, 278
SCHNEEBERGER, E. E., LEBER, P. D., KARNOVSKY, M. J., and MCCLUSKEY, R. T. Altered functional properties of the renal glomerulus in autologous immune complex nephritis. An ultrastructural tracer study, 1283
SCHRADER, JOHN W. Induction of immunological tolerance to a thymus-dependent antigen in the absence of thymus-derived cells, 1303
— and NOSSAL, G. J. V. Effector cell blockade. A new mechanism of immune hyporeactivity induced by multivalent antigens, 1582
SCHUMANN, GEBHARD. See DUKOR, SCHUMANN, GISLER, DIERICH, KÖNIG, HADDING, and BITTER-SUERMANN, 337
SELA, MICHAEL. See FUCHS, MOZES, MAOZ, and SELA, 148
SELL, KENNETH W. See AHMED, STRONG, SELL, THURMAN, KNUDSEN, WISTAR, and GRACE, 902
SHAW, ANDREW, BERKO, BARBARA, and WEGMANN, THOMAS G. Immunological tolerance. Dissociation between in vivo and in vitro reactivity in parabiosed mice, 767
SHEARER, GENE M. See LOZNER, SACHS, and SHEARER, 1204
SHEARER, WILLIAM T., PHILPOTT, GORDON W., and PARKER, CHARLES W. Humoral immunostimulation. I. Increased uptake of [^{125}I]iododeoxyuridine and [^{3}H]thymidine into TNP cells treated with anti-TNP antibody, 367
SHEVACH, ETHAN M., GREEN, IRA, and PAUL, WILLIAM E. Alloantiserum-induced inhibition of immune response gene product function. II. Genetic analysis of target antigens, 679
—, PAUL, WILLIAM E., and GREEN, IRA. Alloantiserum-induced inhibition of immune response gene product function. I. Cellular distribution of target antigens, 661
SHOPE, THOMAS C. See KAPLAN, SHOPE, and PETERSON, 1070
SILVERSTEIN, SAMUEL C. See GRIFFIN and SILVERSTEIN, 323
SLEIGHT, DOUGLAS R. See CHOI, SLEIGHT, and NISONOFF, 761
SLETTEN, K. See HUSBY, NATVIG, and SLETTEN, 773
SNOW, BEVERLEY F. See OATES, STOKES, STOREY, GLOVER, and SNOW, 239
SPEAR, PATRICIA G., and EDELMAN, GERALD M. Maturation of the humoral immune response in mice, 249
SPICER, S. S. See GARVIN, SPICER, PARMLEY, and MUNSTER, 1077
SPIEGELBERG, HANS L. See VALLOTA, GÖTZE, SPIEGELBERG, FORRISTAL, WEST, and MÜLLER-EBERHARD, 1249
SPRENT, J., and MILLER, J. F. A. P. Effect of recent antigen priming on adoptive immune responses. II. Specific unresponsiveness of circulating lymphocytes from mice primed with heterologous erythrocytes, 1

STANKOVA, LIBUSE. See BIGLEY and STANKOVA, 1084
STASHAK, PHILIP W. See AMSBAUGH, HANSEN, PRESCOTT, STASHAK, ASOFSKY, and BAKER, 1499
STEINMAN, RALPH M., and COHN, ZANVIL A. Identification of a novel cell type in peripheral lymphoid organs of mice. II. Functional properties in vitro, 380
—, LUSTIG, DINAH S., and COHN, ZANVIL A. Identification of a novel cell type in peripheral lymphoid organs of mice. III. Functional properties in vivo, 1431
STEJSKAL, V. See O'TOOLE, STEJSKAL, PERLMANN, and KARLSSON, 457
STEPHENSON, JOHN R., TRONICK, STEVEN R., REYNOLDS, ROBERTA K., and AARONSON, STUART A. Isolation and characterization of C-type viral gene products of virus-negative mouse cells, 427
STOCKER, JOHN W., MARCHALONIS, JOHN J., and HARRIS, ALAN W. Inhibition of a T-cell-dependent immune response in vitro by thymoma cell immunoglobulin, 785
STOKES, GORDON S. See OATES, STOKES, STOREY, GLOVER, and SNOW, 239
STOREY, BRIAN G. See OATES, STOKES, STOREY, GLOVER, and SNOW, 239
STREUN, EDWIN W. See RUBINSTEIN, LIU, STREUN, and DECARY, 313
STRONG, DOUGLAS M. See AHMED, STRONG, SELL, THURMAN, KNUDSEN, WISTAR, and GRACE, 902
SUSSMAN, MICHAEL D. See ROWE, MCGOODWIN, MARTIN, SUSSMAN, GRAHN, FARIS, and FRANZBLAU, 180
SUTHERLAND, DAVID E. R. See LEE, MAUER, BROWN, SUTHERLAND, MICHAEL, and NAJARIAN, 793

TADA, TOMIO. See TANIGUCHI and TADA, 108
TANIGUCHI, MASARU, and TADA, TOMIO. Dual regulatory role of the thymus in the maturation of immune response in the rabbit, 108
THEOFILOPOULOS, ARGYRIOS N., BOKISCH, VIKTOR A., and DIXON, FRANK J. Receptor for soluble C3 and C3b on human lymphoblastoid (Raji) cells. Properties and biological significance, 696
THURMAN, GARY B. See AHMED, STRONG, SELL, THURMAN, KNUDSEN, WISTAR, and GRACE, 902
TITTOR, WOLFGANG, GERBASE-DELIMA, MARIA, and WALFORD, ROY L. Synergy among responding lymphoid cells in the one-way mixed lymphocyte reaction. Interaction between two types of thymus-dependent cells, 1488
TODD, JEAN. See GORDON, TODD, and COHN, 1228
TOIVANEN, AULI. See TOIVANEN, TOIVANEN, and VAINIO, 1344
TOIVANEN, PAAVO, TOIVANEN, AULI, and VAINIO, OLLI. Complete restoration of bursa-dependent immune system after transplantation of semiallogeneic stem cells into immunodeficient chicks, 1344
TRAININ, NATHAN. See KOOK and TRAININ, 193
TREFFERS, H. PETER. See HA, WAKSMAN, and TREFFERS, 13
TRONICK, STEVEN R. See STEPHENSON, TRONICK, REYNOLDS, and AARSONSON, 427
TUTHILL, RALPH. See DAHL and TUTHILL, 617

UHR, JONATHAN W., and VITETTA, ELLEN S. Cell surface immunoglobulin. VIII. Synthesis, secretion, and cell surface expression of immunoglobulin in murine thoracic duct lymphocytes, 1013
—. See VITETTA, GRUNDKE-IQBAL, HOLMES, and UHR, 862
—. See VITETTA and UHR, 1599

UNANUE, EMIL R., AULT, KENNETH A., and KARNOVSKY, MORRIS J. Ligand-induced movement of lymphocyte surface macromolecules. IV. Stimulation of cell motility by anti-Ig and lack of relationship to capping, 295
—. See AULT and UNANUE, 1110
UNKELESS, JAY C., GORDON, SAIMON, and REICH, E. Secretion of plasminogen activator by stimulated macrophages, 834
UTERMOHLEN, VIRGINIA, WINFIELD, JOHN B., ZABRISKIE, JOHN B., and KUNKEL, HENRY G. A depression of cell-mediated immunity to measles antigen in patients with systemic lupus erythematosus, 1019

VAINIO, OLLI, See TOIVANEN, TOIVANEN, and VAINIO, 1344
VALLOTA, ENRIQUE H., GÖTZE, OTTO, SPIEGELBERG, HANS L., FORRISTAL, JUDITH, WEST, CLARK D., and MÜLLER-EBERHARD, HANS J. A serum factor in chronic hypocomplementemic nephritis distinct from immunoglobulins and activating the alternate pathway of complement, 1249
VAN BOXEL, JOHN A., and ROSENSTREICH, DAVID L. Binding of aggregated γ-globulin to activated T lymphocytes in the guinea pig, 1002
VAN DEN ENGH, GERRIT, J., and GOLUB, EDWARD S. Antigenic differences between hemopoietic stem cells and myeloid progenitors, 1621
VITETTA, ELLEN S., GRUNDKE-IQBAL, INGE, HOLMES, KATHRYN V., and UHR, JONATHAN W. Cell surface immunoglobulin. VII. Synthesis, shedding, and secretion of immunoglobulin by lymphoid cells of germ-free mice, 862
— and UHR, JONATHAN W. Cell surface immunoglobulin. IX. A new method for the study of synthesis, intracellular transport, and exteriorization in murine splenocytes, 1599
VITETTA, ELLEN S. See UHR and VITETTA, 1013
VOLKERT, MOGENS, MARKER, OLE, and BRO-JØRGENSEN, KLAUS. Two populations of T lymphocytes immune to the lymphocytic choriomeningitis virus, 1329
VOLKMAN, ALVIN, and COLLINS, FRANK M. The cytokinetics of monocytosis in acute *Salmonella* infection in the rat, 264

WAKSMAN, BYRON H. See HA, WAKSMAN, and TREFFERS, 13
WALFORD, ROY L. See TITTOR, GERBASE-DELIMA, and WALFORD, 1488
WECKER, E. See HÜNIG, SCHIMPL, and WECKER, 754
WEGMANN, THOMAS G. See SHAW, BERKO, and WEGMANN, 767
WEIGERT, MARTIN, RASCHKE, WILLIAM C., CARSON, DENNIS, and COHN, MELVIN. Immunochemical analysis of the idiotypes of mouse myeloma proteins with specificity for levan or dextran, 137
WEIGLE, WILLIAM O. See CLAGETT and WEIGLE, 643
—. See LAWRENCE and WEIGLE, 943
WEILER, EBERHARDT. See KOLB, DI PAULI, and WEILER, 467
WEST, CLARK D. See VALLOTA, GÖTZE, SPIEGELBERG, FORRISTAL, WEST, and MÜLLER-EBERHARD, 1249
WIGZELL, HANS. See BINZ, LINDENMANN, and WIGZELL, 877
—. See RUBIN and WIGZELL, 732
—. See SCHIRRMACHER, RUBIN, PROSS, and WIGZELL, 93
WILLENBORG, DAVID O., and PRENDERGAST, ROBERT A. The effect of sea star coelomocyte extract on cell-mediated resistance to *Listeria monocytogenes* in mice, 820

WILLIAMS, ALAN F. See HUNT and WILLIAMS, 479
WILLIAMSON, A. R. See MCMICHAEL and WILLIAMSON, 1361
WINCHESTER, ROBERT J. See FU, WINCHESTER, and KUNKEL, 451
—. See KUNKEL, WINCHESTER, JOSLIN, and CAPRA, 128
WINFIELD, JOHN B. See UTERMOHLEN, WINFIELD, ZABRISKIE, and KUNKEL, 1019
WISTAR, RICHARD, JR. See AHMED, STRONG, SELL, THURMAN, KNUDSEN, WISTAR, and GRACE, 902
WOLFF, SHELDON M. See DINARELLO, GOLDIN, and WOLFF, 1369

YOSHIKI, TAKASHI, MELLORS, ROBERT C., HARDY, WILLIAM D., JR., and FLEISSNER, ERWIN. Common cell surface antigen associated with mammalian C-type RNA viruses. Cell membrane-bound gs antigen, 925

ZABRISKIE, JOHN B. See UTERMOHLEN, WINFIELD, ZABRISKIE, and KUNKEL, 1019

INDEX TO SUBJECTS

ACID hydrolases, lysosomal, extracellular release in macrophages exposed to group A streptococcal cell wall substance, and changes in cellular enzyme levels, 1262

—, RNA, tumor virus, endogenous, autogenous immunity to, identification of antibody reactivity to select viral antigens, 1568

—, RNA, viruses, mammalian C-type, common cell surface antigen associated with, 925

Agglutinability and electrophoretic mobility of red blood cells, polymorphism in mice, 313

Alloantibodies, characterization of receptor-carrying cells by use of antibodies to, 877

Alloantigens, cell-bound receptors for on normal lymphocytes, 877

Alloantiserum-induced inhibition of immune response gene product function, cellular distribution of target antigens, 661

—inhibition of immune response gene product function, genetic analysis of target antigens, 679

Allogeneic cell interactions and enzymatic digestion with trypsin, effects on inactivated hapten-specific precursors of antibody-forming cells, 1446

— lymphocytes and phytomitogens, cellular interactions in proliferative response of human T and B lymphocytes to, 1553

— or co-isogenic thymuses, activity of host-derived T cells which differentiate in nude mice grafted with, 1215

Allogeneically stimulated human lymphocyte cell lines, partial purification of nonspecific enhancing factor from supernates of, 439

Allograft, cell-mediated, and humoral antibody reactions, induction, 398

Allografts, bone marrow, role of ^{89}Sr-sensitive effector cells responsible for rejection of, 1093

Allotype, immunoglobulin, commitment of gene expression restriction in B lymphocytes and progeny to, 581

—, immunoglobulin, hidden, expression in congenic partner strain of BALB/c mice, 512

Amyloid fibril protein, new, third class, 773

Angiotensin I and II, renal hypertension in rats immunized against, 239

Antibodies, idiotypic, production and characterization of autoantiidiotypic antisera, 712

—, restricted, to streptococcal group polysaccharides in mice, genetics, strain differences of isoelectric focusing spectra of group A hyperimmune antisera, 1513

— to alloantibodies, characterization of receptor-carrying cells by use of, 877

Antibody, anti-TNP, increased uptake of [^{125}I]iododeoxyuridine and [^{3}H]thymidine into TNP cells treated with, 367

— as mediator of specific cytotoxicity, 93

— formation, potentiation of T-cell-mediated immunity by selective suppression of with cyclophosphamide, 1529

—, humoral, and cell-mediated allograft reactions, induction, 398

—, natural, to phosphorylcholine, genetics of new IgV_H(T15 idiotype) marker in mouse regulating, 983

Antibody, production to hapten 2,4-dinitrobenzene, stimulation by affinity labeled murine lymphoid cells, 943
— reactivity to viral antigens, identification of, autogenous immunity to endogenous RNA tumor virus, 1568
—, receptor, anti-T-cell, inhibition of specific cell-mediated cytotoxicity by, 888
— response, primary, in vitro, to DNP-Ficoll, cellular requirements for, 1354
— response to nuclease by *Ir* region of mouse *H-2* complex, control of, 1204
— response to Type III pneumococcal polysaccharide in mice, genetic control, relationship between IgM immunoglobulin levels and ability to give IgM antibody response, 1499
Antibody-forming cells, effects of allogeneic cell interactions and enzymatic digestion with trypsin on inactivated hapten-specific precursors of, 1446
Antibody-hapten complexes, persistence in circulation of immunized animals after single intravenous hapten injection, 278
Antibody-induced movement of membrane components of *Leishmania enrietii*, 1061
Antibody-producing cells, autoradiographic studies on proliferation in vitro, 754
Antigen binding to lymphocytes, removal and regeneration of receptor, 1110
—, cell surface, common, associated with mammalian C-type RNA viruses, 925
— injection, dose and route, influence on immunological induction of T cells, 528
—, measles, depression of cell-mediated immunity to in patients with systemic lupus erythematosus, 1019
— priming, recent, effect on adoptive immune responses, 1
Antigen, SV40 T and human chromosome 7, segregation of expression in mouse-human hybrid subclones, 1350
—, thymus-dependent, induction of immunological tolerance to in absence of thymus-derived cells, 1303
Antigenic differences between hemopoietic stem cells and myeloid progenitors, 1621
Antigenicity, cellular, progressive loss in Gross virus lymphoma, and induction of partial immunologic tolerance in rats, 1382
Antigens, HL-A, and $\beta 2$-microglobulin, relationship between studied by immunofluorescence on lymphocyte membrane, 957
—, multivalent, new mechanism of immune hyporeactivity induced by, 1582
—, soluble, defined, cytotoxic immune cells with specificity for, 93
—, target, cellular distribution in alloantiserum-induced inhibition of immune response gene product function, 661
—, target, genetic analysis in alloantiserum-induced inhibition of immune response gene product function, 679
—, viral, identification of antibody reactivity to, autogenous immunity to endogenous RNA tumor virus, 1568
Antisera, autoantiidiotypic, production and characterization, 712
—, hyperimmune, group A, strain differences of isoelectric focusing spectra of, genetics of restricted antibodies to streptococcal group polysaccharides in mice, 1513
Ascorbate, oxidized and reduced, uptake and reduction by human leukocytes, 1084
Autoantiidiotypic antisera, production and characterization, 712
Autoimmune-like activity in syngeneic and semisyngeneic systems, 230

Autoradiographic studies on proliferation of antibody-producing cells in vitro, 754

BACILLUS Calmette-Guérin, modifying effect on immunological induction of T cells, 1540
Bacterial lipopolysaccharide, regulatory influence on specific T-cell helper function, 24
Basophilic leukemia cells and IgE, interaction, evidence for specific binding of IgE, 600
Besnoitia and *Toxoplasma* in specifically and cross-immunized hamsters and in cultures, cell-mediated immunity against, 560
Bladder, urinary, carcinoma, lymphoid cells mediating tumor-specific cytotoxicity to, separation of effector population using surface marker, 457
Blood cells, red. *See* Erythrocytes.
— pressure, increased, and mortality in spontaneously hypertensive rat, 617
Bone marrow allografts, role of ^{89}Sr-sensitive effector cells responsible for rejection of, 1093
— marrow and thymus cell cooperation in immune response to gelatin, thymus independence of collagen-like synthetic polypeptide and of collagen and need for, 148
— marrow-derived cell, complement-dependent activation by cobra venom factor and other mitogens, 337
— marrow-derived cell mitogens, effect on lymphocyte subpopulations possessing C′3 and Fc receptors, 969
— marrow-derived cells bearing specific receptors, method for isolation and recovery, 761
— marrow-derived cells, mitogenic activation results in specific immune responses, 74
— marrow-derived lymphocytes, immunological tolerance in, 1446, 1464
Bone marrow-derived lymphocytes, ontogeny, 1125, 1142
— marrow-derived lymphocytes and progeny, gene expression restriction in, commitment to immunoglobulin allotype, 581
— marrow-derived memory cells, time-course of proliferation of, 1361
— marrow-derived and thymus-derived human lymphocytes, cellular interactions in proliferative response to phytomitogens and allogeneic lymphocytes, 1553
— marrow-derived and thymus-derived lymphocytes, activation in vitro, 24
— marrow-derived and thymus-derived lymphocytes, roles in termination of unresponsiveness to autologous thyroglobulin, 643
— marrow-derived and thymus-derived lymphocytes and thymocytes, presence of receptors for aggregated IgG on, 1175
Brain and thymus lipid inhibition of antibrain-associated θ-cytotoxicity, 445
Bursa-dependent immune system, complete restoration after transplantation of semiallogeneic stem cells into immunodeficient chicks, 1344

CALMETTE-GUÉRIN bacillus, modifying effect on immunological induction of T cells, 1540
Carcinoma, urinary bladder, lymphoid cells mediating tumor-specific cytotoxicity to, separation of effector population using surface marker, 457
Carrier-hapten helper effect and cell-mediated immunity to carrier, lack of correlation between, 779
Cells, blood, red. *See* Erythrocytes.
Cellular antigenicity, progressive loss in Gross virus lymphoma, and induction of partial immunologic tolerance in rats, 1382
— distribution of target antigens in alloantiserum-induced inhibition of

immune response gene product function, 661
Cellular enzyme levels, changes in, and extracellular release of lysosomal acid hydrolases in macrophages exposed to group A streptococcal cell wall substance, 1262
— immunity, mediator, assembly of sensitized lymphocytes in inflammatory exudates, 1415
— interactions in proliferative response of human T and B lymphocytes to phytomitogens and allogeneic lymphocytes, 1553
— requirements for primary in vitro antibody response to DNP-Ficoll, 1354
— stimulation, lectin-induced, interaction of cholera toxin and toxin derivatives with lymphocytes, binding properties and interference with, 801
Chains, K-, of immunoglobulin from continuous culture of human lymphocytes, Daudi, have unusual molecular size, 1031
—, light, of anti-γ-globulins, showing cross-idiotypic specificities, similarities in, 128
—, light, surface IgM, IgD, and free, on human lymphocytes, 451
Cholera toxin and toxin derivatives, interaction with lymphocytes, binding properties and interference with lectin-induced cellular stimulation, 801
Choriomeningitis virus, lymphocytic, two populations of T lymphocytes immune to, 1329
Chromosome 7, human, and SV40 T antigen, segregation of expression in mouse-human hybrid subclones, 1350
Circulation of immunized animals, persistence of hapten-antibody complexes after single intravenous hapten injection, 278
Clonal memory, time-course of proliferation of B-memory cells, 1361
— nature of immune response to phosphorylcholine, 58
Cobra venom factor and other mitogens, complement-dependent B-cell activation by, 337
Coelomocyte, sea star, extract, effect on cell-mediated resistance to *Listeria monocytogenes*, 820
Collagen and elastin, sex-linked defect in cross-linking associated with mottled locus in mice, 180
Collagen-like synthetic polypeptide and collagen, thymus independence of, need for thymus and bone marrow cell cooperation in immune response to gelatin, 148
Complement activation, properdin role in alternate pathway of, 44
— pathway, alternate, serum factor in chronic hypocomplementemic nephritis distinct from immunoglobulins and activating, 1249
— receptor lymphocytes, *H-2* linkage of gene controlling rate of appearance of, 1142
— receptors and surface immunoglobulin, relative rates of appearance of lymphocytes bearing, 1125
Complement-dependent B-cell activation by cobra venom factor and other mitogens, 337
Conjugates, hapten, of unrelated immunogenic or nonimmunogenic carriers, tolerance induction in prime B cells by, 1464
—, protein, hapten autologous, lymphocyte proliferation in vitro induced by, 732
Cyclophosphamide, potentiation of T-cell-mediated immunity by selective suppression of antibody formation with, 1529
Cytokinetics of monocytosis in acute *Salmonella* infection, 264
Cytotoxic activity of mouse macrophages studied by inhibitors, 1049
— immune cells with specificity for defined soluble antigens, 93
θ-Cytotoxicity, antibrain-associated, brain and thymus lipid inhibition of, 445

Cytotoxicity, cell-mediated, specific, inhibition by anti-T-cell receptor antibody, 888
—, tumor-specific, to urinary bladder carcinoma, lymphoid cells mediating, separation of effector population using surface marker, 457

DEXTRAN or levan, immunochemical analysis of idiotypes of myeloma proteins with specificity for, 137
Dextrans, immunochemical studies on myeloma proteins reactive with, or with fructosans and on human antilevans, 159
Diabetes mellitus in rats, renal transplantation in, 793
— mellitus-like disease in mice infected with encephalomyocarditis virus, genetic influences affecting occurrence of, 414
Digestion, enzymatic, with trypsin and allogeneic cell interactions, effects on inactivated hapten-specific precursors of antibody-forming cells, 1446
2,4-Dinitrobenzene, hapten, stimulation of antibody production to by affinity labeled murine lymphoid cells, 943
Dinitrophenyl-Ficoll, cellular requirements for primary in vitro antibody response to, 1354
Disease, diabetes mellitus-like, in mice infected with encephalomyocarditis virus, genetic influences affecting occurrence of, 414
—, Hodgkin's, immunohistochemical demonstration of IgG in Reed-Sternberg and other cells in, 1077
Dose and route of antigen injection, influence on immunological induction of T cells, 528
Duct, thoracic, murine, lymphocytes, synthesis, secretion, and cell surface expression of immunoglobulin in, 1013

ELASTIN and collagen, sex-linked defect in cross-linking associated with mottled locus in mice, 180
Electron, immuno-, microscope tracer consisting of heme-octapeptide coupled to Fab, preparation and characterization, 208
Electrophoretic mobility and agglutinability of red blood cells, new polymorphism in mice, 313
Encephalomyocarditis virus, genetic influences affecting occurrence of diabetes mellitus-like disease in mice infected with, 414
Enteric infection, route in normal mice, 1189
Enzymatic digestion with trypsin and allogeneic cell interactions, effects on inactivated hapten-specific precursors of antibody-forming cells, 1446
Enzyme levels, cellular, changes in, and extracellular release of lysosomal acid hydrolases in macrophages exposed to group A streptococcal cell wall substance, 1262
— release, lysosomal, from human neutrophils, hormonal control, effects of autonomic agents on enzyme release, phagocytosis, and cyclic nucleotide levels, 1395
Epithelial cells, intestinal, human, localization of free and bound secretory component in, model for assembly of secretory IgA, 629
Epstein-Barr virus-negative human malignant T-cell lines, 1070
Erythrocytes, electrophoretic mobility and agglutinability, new polymorphism in mice, 313
—, heterologous, specific unresponsiveness of circulating lymphocytes from mice primed with, 1
Exudates, inflammatory, assembly of sensitized lymphocytes in, 1415

FIBRIL protein, amyloid, new, third class, 773
Fluid, spinal, and plasma of patients with subacute sclerosing panen-

cephalitis, partial characterization of blocking factor in, 902
Friend virus leukemia in mice, mechanisms of resistance, 1093
Fructosans, immunochemical studies on myeloma proteins reactive with dextrans or with, and on human antilevan, 159

GELATIN, thymus independence of collagen-like synthetic polypeptide and of collagen, need for thymus and bone marrow cell cooperation in immune response to, 148
Gene controlling rate of appearance of complement receptor lymphocytes, *H-2* linkage, 1142
— expression, restriction in B lymphocytes and progeny, commitment to immunoglobulin allotype, 581
— product function, immune response, alloantiserum-induced inhibition of, cellular distribution of target antigens, 661
— product function, immune response, alloantiserum-induced inhibition of, genetic analysis of target antigens, 679
— products, C-type viral, of virus-negative mouse cells, isolation and characterization, 427
Genetic analysis of responses to *H-2D*^{d} specificities, in mixed lymphocyte culture responses of mice, 851
— analysis of target antigens in alloantiserum-induced inhibition of immune response gene product function, 679
— control of antibody response to Type III pneumococcal polysaccharide in mice, relationship between IgM immunoglobulin levels and ability to give IgM antibody response, 1499
— control of immune response to staphylococcal nuclease, 1204
— influences affecting occurrence of diabetes mellitus-like disease in mice infected with encephalomyocarditis virus, 414
Genetic resistance to Friend virus leukemia in mice, mechanisms, 1093
Genetics of new IgV_H(T15 idiotype) marker in mouse regulating natural antibody to phosphorylcholine, 983
— of restricted antibodies to streptococcal group polysaccharides in mice, strain differences of isoelectric focusing spectra of group A hyperimmune antisera, 1513
Germ-free mice, synthesis, shedding, and secretion of immunoglobulin in lymphoid cells of, 862
γ-Globulin, aggregated, binding to activated T lymphocytes, 1002
γ-Globulins, anti-, showing cross-idiotypic specificities, similarities in light chains of, 128
Glomerulus, renal, altered functional properties in autologous immune complex nephritis, ultrastructural tracer study, 1283
Graft-vs.-host assays, two, dissociation of T-cell activities in mixed lymphocyte reaction and, 721
Gross virus lymphoma, induction of partial immunologic tolerance in rats and progressive loss of cellular antigenicity in, 1382

HAPTEN autologous protein conjugates, lymphocyte proliferation in vitro induced by, 732
— conjugates of unrelated immunogenic or nonimmunogenic carriers, tolerance induction in primed B cells by, 1464
— 2,4-dinitrobenzene, stimulation of antibody production to by affinity labeled murine lymphoid cells, 943
Hapten-antibody complexes, persistence in circulation of immunized animals after single intravenous hapten injection, 278
Hapten-carrier helper effect and cell-mediated immunity to carrier, lack of correlation between, 779
Hapten-specific, inactivated, precursors of antibody-forming cells,

effects of allogeneic cell interactions and enzymatic digestion with trypsin on, 1446
Heme-octapeptide coupled to Fab, preparation and characterization of immunoelectron microscope tracer consisting of, 208
Hemopoietic stem cells and myeloid progenitors, antigenic differences between, 1621
Heterologous enhancement, specific, of immune responses, 439
— erythrocytes, specific unresponsiveness of circulating lymphocytes from mice primed with, 1
Histochemical, immuno-, demonstration of IgG in Reed-Sternberg and other cells in Hodgkin's disease, 1077
Hodgkin's disease, immunohistochemical demonstration of IgG in Reed-Sternberg and other cells in, 1077
Hormonal control of lysosomal enzyme release from human neutrophils, effects of autonomic agents on enzyme release, phagocytosis, and cyclic nucleotide levels, 1395
Hormone-like activity of thymus humoral factor on induction of immune competence in lymphoid cells, 193
Host-derived T cells which differentiate in nude mice grafted with coisogenic or allogeneic thymuses, activity of, 1215
Host-vs.-graft assays, two, dissociation of T-cell activities in mixed lymphocyte reaction and, 721
Humoral antibody and cell-mediated allograft reactions, induction, 398
— factor, thymus, hormone-like activity on induction of immune competence in lymphoid cells, 193
— immune response, maturation, 249
— immunostimulation, increased uptake of [^{125}I]iododeoxyuridine and [^{3}H]thymidine into TNP cells treated with anti-TNP antibody, 367
Hybrid mice, reciprocal, graft-vs.-host reactions in, 721
— subclones, mouse-human, segregation of expression of SV40 T antigen and human chromosome 7 in, 1350
Hydrolases, acid, lysosomal, extracellular release in macrophages exposed to group A streptococcal cell wall substance, and changes in cellular enzyme levels, 1262
Hyperimmune antisera, group A, strain differences of isoelectric focusing spectra of, genetics of restricted antibodies to streptococcal group polysaccharides in mice, 1513
Hypertension, renal, in rats immunized against angiotensin I and II, 239
Hypertensive, spontaneously, rat, increased blood pressure and mortality in, 617
Hypocomplementemic nephritis, chronic, serum factor distinct from immunoglobulins and activating alternate complement pathway in, 1249
Hyporeactivity, immune, induced by multivalent antigens, new mechanism of, 1582

IDIOTYPE, specificity, and class of phosphorylcholine-binding receptors on lymphoid cells, clonal nature of immune response, 58
—, T15, new IgV_H marker in mouse regulating natural antibody to phosphorylcholine, genetics of, 983
Idiotypes of myeloma proteins with specificity for levan or dextran, immunochemical analysis of, 137
Idiotypic antibodies, production and characterization of autoantiidiotypic antisera, 712
—, cross-, specificities, similarities in light chains of anti-γ-globulins showing, 128
Immunity, autogenous, to endogenous RNA tumor virus, identification of

antibody reactivity to select viral antigens, 1568
Immunity, cell-mediated, against *Besnoitia* and *Toxoplasma* in specifically and cross-immunized hamsters and in cultures, 560
—, cell-mediated, to measles antigen, depression in patients with systemic lupus erythematosus, 1019
—, cellular, mediator, assembly of sensitized lymphocytes in inflammatory exudates, 1415
—, T-cell-mediated, potentiation by selective suppression of antibody formation with cyclophosphamide, 1529
Immunization against angiotensin I and II, renal hypertension in rats, 239
Immunized animals, persistence of hapten-antibody complexes in circulation after single intravenous hapten injection, 278
Immunochemical analysis of idiotypes of myeloma proteins with specificity for levan or dextran, 137
— studies on myeloma proteins reactive with dextrans or with fructosans and on human antilevans, 159
Immunocyte triggering, thymus-independent, mitogenic activation of B cells results in specific immune responses, 74
Immunodeficient chicks, complete restoration of bursa-dependent immune system after transplantation of semiallogeneic stem cells into, 1344
Immunoelectron microscope tracer consisting of heme-octapeptide coupled to Fab, preparation and characterization, 208
Immunofluorescence, relationship between HL-A antigens and β2-microglobulin studied by on lymphocyte membrane, 957
Immunogenic, unrelated, or nonimmunogenic carriers, tolerance induction in primed B cells by hapten conjugates of, 1464
Immunoglobulin A, secretory, model for assembly, 629
— allotype, commitment of gene expression restriction in B lymphocytes and progeny to, 581
— allotype, hidden, expression in congenic partner strain of BALB/c mice, 512
—, cell surface, new method for study of synthesis, intracellular transport, and exteriorization in murine splenocytes, 1599
—, cell surface, of marrow-derived and thymus-derived lymphocytes, origin, 479
—, cell surface, synthesis, secretion, and expression in murine thoracic duct lymphocytes, 1013
—, cell surface, synthesis, shedding, and secretion by lymphoid cells of germ-free mice, 862
— E and basophilic leukemia cells, interaction, evidence for specific binding of IgE, 600
— from continuous culture of human lymphocytes, Daudi, K-chains have unusual molecular size, 1031
— G, aggregates, receptors for on mouse lymphocytes, 1175
— G, immunohistochemical demonstration in Reed-Sternberg and other cells in Hodgkin's disease, 1077
— G, induction by lipid A in newborn mouse, 467
—, IgM, levels, and ability to give IgM antibody response, relationship between, 1499
— M, immunoglobulin D, and free light chains, surface, occurrence on human lymphocytes, 451
—, surface, and complement receptors, relative rates of appearance of lymphocytes bearing, 1125
—, thymoma cell, inhibition of T-cell-dependent immune response in vitro by, 785
Immunoglobulins, serum factor in chronic hypocomplementemic ne-

phritis distinct from and activating alternate complement pathway, 1249
Immunohistochemical demonstration of IgG in Reed-Sternberg and other cells in Hodgkin's disease, 1077
Immunologic tolerance, partial, induction in rats and progressive loss of cellular antigenicity in Gross virus lymphoma, 1382
Immunological induction of T cells, influence of dose and route of antigen injection on, 528
— induction of T cells, modifying effect of BCG on, 1540
— tolerance, dissociation between in vivo and in vitro reactivity in parabiosed mice, 767
— tolerance in bone marrow-derived lymphocytes, 1446, 1464
— tolerance to thymus-dependent antigen, induction in absence of thymus-derived cells, 1303
Immunostimulation, humoral, increased uptake of [^{125}I]iododeoxyuridine and [^{3}H]thymidine into TNP-cells treated with anti-TNP antibody, 367
Inflammatory exudates, assembly of sensitized lymphocytes in, 1415
Inhibition, alloantiserum-induced, of immune response gene product function, cellular distribution of target antigens, 661
—, alloantiserum-induced, of immune response gene product function, genetic analysis of target antigens, 679
—, brain and thymus lipid, of anti-brain-associated θ-cytotoxicity, 445
—, feedback, of specifically sensitized lymphocytes, 543
— of specific cell-mediated cytotoxicity by anti-T-cell receptor antibody, 888
— of T-cell-dependent immune response in vitro by thymoma cell immunoglobulin, 785
Inhibitors, cytotoxic activity of mouse macrophages studied by, 1049
Intestinal epithelial cells, human, localization of free and bound secretory component in, model for assembly of secretory IgA, 629
Intracellular transport, synthesis, and exteriorization of cell surface immunoglobulin in murine splenocytes, new method for study, 1599
Iododeoxyuridine, ^{125}I-labeled, and thymidine, ^{3}H-labeled, increased uptake into TNP cells treated with anti-TNP antibody, 367

KIDNEY glomerulus, altered functional properties in autologous immune complex nephritis, ultrastructural tracer study, 1283
Kidney hypertension in rats immunized against angiotensin I and II, 239
— transplantation in diabetes mellitus in rats, 793
Kinetics, cell cycle, and responder recruitment, stimulated lymphocyte cultures, 1037

LECTIN-INDUCED cellular stimulation, interaction of cholera toxin and toxin derivatives with lymphocytes, binding properties and interference with, 801
Leishmania enrietii, antibody-induced movement of membrane components of, 1061
Leukemia cells, basophilic, and IgE, interaction, evidence for specific binding of IgE, 600
— virus, Friend, in mice, mechanisms of genetic resistance to, 1093
Leukocyte, mixed, stimulatory, M-locus product, absence from thymus cells, 1628
Leukocytes, human, uptake and reduction of oxidized and reduced ascorbate by, 1084
Leukocytic pyrogens, human, two, demonstration and characterization, 1369
Levan or dextran, immunochemical

analysis of idiotypes of myeloma proteins with specificity for, 137
Ligand-induced movement of lymphocyte surface macromolecules, stimulation of cell motility by anti-Ig and lack of relationship to capping, 295
Lipid A induction of IgG in newborn mouse, 467
—, thymus, and brain inhibition of antibrain-associated θ-cytotoxicity, 445
Lipopolysaccharide, bacterial, regulatory influence on specific T-cell helper function, 24
— is thymus dependent, Peyer's patch lymphoid cells, 407
Listeria monocytogenes, effect of sea star coelomocyte extract on cell-mediated resistance to, 820
Lupus erythematosus, systemic, depression of cell-mediated immunity to measles antigen in patients with, 1019
Lymphoblastoid, Raji, cells, human, receptor for soluble C3 and C3b on, properties and biological significance, 696
Lymphocyte cell lines, human, allogeneically stimulated, partial purification of nonspecific enhancing factor from supernates of, 439
— culture responses, mixed, of mice, genetic analysis of responses to $H\text{-}2D^d$ specificities, 851
— cultures, murine mixed, effect of 2-mercaptoethanol on, 1025
— cultures, stimulated, responder recruitment and cell cycle kinetics, 1037
— membrane, relationship between HL-A antigens and β2-microglobulin studied by immunofluorescence on, 957
— proliferation in vitro induced by hapten autologous protein conjugates, 732
— reaction, mixed, one-way, synergy among responding lymphoid cells in, interaction between two types of thymus-dependent cells, 1488
Lymphocyte reaction, mixed, and two graft-vs.-host assays, dissociation of T-cell activities in, 721
— reactivity, mixed, against normal cells by splenic lymphocytes from tumor-bearing mice, 224, 230
— subpopulations possessing C′3 and Fc receptors, effect of B-cell mitogens on, 969
— surface macromolecules, ligand-induced movement, stimulation of cell motility by anti-Ig and lack of relationship to capping, 295
Lymphocytes, antigen binding to, removal and regeneration of receptor, 1110
—, B, ontogeny, 1125, 1142
—, B, and progeny, restriction of gene expression in, commitment to immunoglobulin allotype, 581
—, bone marrow-derived, immunological tolerance in, 1446, 1464
—, circulating, from mice primed with heterologous erythrocytes, specific unresponsiveness, 1
—, human, Daudi, continuous culture, K-chains of immunoglobulin have unusual molecular size, 1031
—, human, occurrence of surface IgM, IgD, and free light chains on, 451
—, interaction of cholera toxin and toxin derivatives with, binding properties and interference with lectin-induced cellular stimulation, 801
—, marrow-derived and thymus-derived, origin of cell surface immunoglobulin of, 479
—, normal, cell-bound receptors for alloantigens on, 877
—, sensitized, assembly in inflammatory exudates, 1415
—, specifically sensitized, feedback inhibition of, 543
—, T, activated, binding of aggregated γ-globulin to, 1002
—, T and B, activation in vitro, 24

Lymphocytes, T and B, human, cellular interactions in proliferative response to phytomitogens and allogeneic lymphocytes, 1553
—, T and B, roles in termination of unresponsiveness to autologous thyroglobulin, 643
—, T, two populations immune to lymphocytic choriomeningitis virus, 1329
—, thoracic duct, murine, synthesis, secretion, and cell surface expression of immunoglobulin in, 1013
—, thymus-derived and bone marrow-derived and thymocytes, presence of receptors for aggregated IgG on, 1175
Lymphocytic choriomeningitis virus, two populations of T lymphocytes immune to, 1329
Lymphoid cells, hormone-like activity of thymus humoral factor on induction of immune competence in, 193
— cells mediating tumor-specific cytotoxicity to urinary bladder carcinoma, separation of effector population using surface marker, 457
— cells, murine, affinity labeled, ability to activate in vivo immune response, 943
— cells of germ-free mice, synthesis, shedding, and secretion of immunoglobulin by, 862
— cells, Peyer's patch, induction of humoral antibody and cell-mediated allograft reactions, 398
— cells, Peyer's patch, lipopolysaccharide is thymus dependent, 407
— cells, specificity, class, and idiotype of phosphorylcholine-binding receptors on, clonal nature of immune response, 58
— cells, synergy among in one-way mixed lymphocyte reaction, interaction between two types of thymus-dependent cells, 1488
— organs, peripheral, identification of novel cell type in, functional properties in vitro, 380
Lymphoid organs, peripheral, identification of novel cell type in, functional properties in vivo, 1431
Lymphoma, Gross virus, induction of partial immunologic tolerance in rats and progressive loss of cellular antigenicity in, 1382
Lysosomal acid hydrolases, extracellular release in macrophages exposed to group A streptococcal cell wall substance, and changes in cellular enzyme levels, 1262
— enzyme release from human neutrophils, hormonal control, effects of autonomic agents on enzyme release, phagocytosis, and cyclic nucleotide levels, 1395
Lysozyme, in vitro synthesis and secretion by mononuclear phagocytes, 1228

MACROMOLECULES, lymphocyte surface, ligand-induced movement of, stimulation of cell motility by anti-Ig and lack of relationship to capping, 295
Macrophage plasma membrane, segmental response to phagocytic stimulus, 323
Macrophages exposed to group A streptococcal cell wall substance, changes in cellular enzyme levels and extracellular release of lysosomal acid hydrolases in, 1262
—, human, normal and activated, effect on *Toxoplasma gondii*, 1154
—, mouse, cytotoxic activity studied by inhibitors, 1049
—, stimulated, secretion of plasminogen activator by, 834
Mammalian C-type RNA viruses, common cell surface antigen associated with, 925
Man, antilevans, immunochemical studies on myeloma proteins reactive with dexrans or with fructosans and on, 159
—, cell cultures, neoplastic, properties of plasminogen activators formed by, 1317
—, chromosome 7, and SV40 T anti-

gen, segregation of expression in mouse-human hybrid subclones, 1350
Man, intestinal epithelial cells, localization of free and bound secretory component in, model for assembly of secretory IgA, 629
—, leukocytes, uptake and reduction of oxidized and reduced ascorbate by, 1084
—, leukocytic pyrogens, two, demonstration and characterization, 1369
—, lymphoblastoid, Raji, cells, receptor for soluble C3 and C3b on, properties and biological significance, 696
—, lymphocyte cell lines, allogeneically stimulated, partial purification of nonspecific enhancing factor from supernates of, 439
—, lymphocytes, Daudi, continuous culture, K-chains of immunoglobulin have unusual molecular size, 1031
—, lymphocytes, occurrence of surface IgM, IgD, and free light chains on, 451
—, macrophages, normal and activated, effect on *Toxoplasma gondii*, 1154
—, malignant T-cell lines, Epstein-Barr virus-negative, 1070
—, neutrophils, hormonal control of lysosomal enzyme release from, effects of autonomic agents on enzyme release, phagocytosis, and cyclic nucleotide levels, 1395
—, protein, acute phase, found in association with rubella virus infection, characterization, 497
—, T and B lymphocytes, cellular interactions in proliferative response to phytomitogens and allogeneic lymphocytes, 1553
Marrow, bone, allografts, role of ^{89}Sr-sensitive effector cells responsible for rejection of, 1093
—, bone, and thymus cell cooperation in immune response to gelatin, thymus independence of collagen-like synthetic polypeptide and of collagen and need for, 148
Marrow-derived, bone, cell mitogens, effect on lymphocyte subpopulations possessing C'3 and Fc receptors, 969
—, bone, cells bearing specific receptors, general method for isolation and recovery, 761
—, bone, cells, mitogenic activation results in specific immune responses, 74
—, bone, lymphocytes, immunological tolerance in, 1446, 1464
—, bone, lymphocytes, ontogeny, 1125, 1142
—, bone, lymphocytes and progeny, gene expression restriction in, commitment to immunoglobulin allotype, 581
—, bone, memory cells, time-course of proliferation of, 1361
—, bone, and thymus-derived human lymphocytes, cellular interactions in proliferative response to phytomitogens and allogeneic lymphocytes, 1553
—, bone, and thymus-derived lymphocytes, activation in vitro, 24
—, bone, and thymus-derived lymphocytes, roles in termination of unresponsiveness to autologous thyroglobulin, 643
—, bone, and thymus-derived lymphocytes and thymocytes, presence of receptors for aggregated IgG on, 1175
— cell, bone, complement-dependent activation by cobra venom factor and other mitogens, 337
— and thymus-derived lymphocytes, origin of cell surface immunoglobulin of, 479
Maturation of humoral immune response, 249
— of immune response, dual regulatory role of thymus in, 108
Measles antigen, depression of cell-mediated immunity to in patients with systemic lupus erythematosus, 1019

Membrane components of *Leishmania enrietii*, antibody-induced movement of, 1061
—, lymphocyte, relationship between HL-A antigens and β2-microglobulin studied by immunofluorescence on, 957
—, macrophage plasma, segmental response to phagocytic stimulus, 323
Membrane-bound, cell, gs antigen associated with mammalian C-type RNA viruses, 925
Memory, clonal, time-course of proliferation of B-memory cells, 1361
2-Mercaptoethanol, effect on murine mixed lymphocyte cultures, 1025
β2-Microglobulin and HL-A antigens, relationship between studied by immunofluorescence on lymphocyte membrane, 957
Microscope tracer, immunoelectron, consisting of heme-octapeptide coupled to Fab, preparation and characterization, 208
Mitogenic activation of B cells results in specific immune responses, thymus-independent immunocyte triggering, 74
Mitogens, B-cell, effect on lymphocyte subpopulations possessing C'3 and Fc receptors, 969
— and cobra venom factor, complement-dependent B-cell activation by, 337
Molecular size, unusual, K-chains of immunoglobulin from continuous culture of human lymphocytes, Daudi, have, 1031
Moloney sarcoma virus, evidence of suppressor cell activity in mice spleens bearing primary tumors induced by, 1473
Monocytosis, cytokinetics, in acute *Salmonella* infection, 264
Mononuclear phagocytes, in vitro synthesis and secretion of lysozyme by, 1228
Mortality and increased blood pressure in spontaneously hypertensive rat, 617
Mottled locus in mice, sex-linked defect in cross-linking of collagen and elastin associated with, 180
Myeloid progenitors and hemopoietic stem cells, antigenic differences between, 1621
Myeloma proteins, immunochemical analysis of idiotypes with specificity for levan or dextran, 137
— proteins reactive with dextrans or with fructosan and on human antilevans, immunochemical studies on, 159

NEOPLASTIC human cell cultures, properties of plasminogen activators formed by, 1317
Nephritis, autologous immune complex, altered functional properties of renal glomerulus in, ultrastructural tracer study, 1283
Nephritis, hypocomplementemic, chronic, serum factor distinct from immunoglobulins and activating complement pathway in, 1249
Neutrophils, human, hormonal control of lysosomal enzyme release from, effects of autonomic agents on enzyme release, phagocytosis, and cyclic nucleotide levels, 1395
Nuclease, staphylococcal, genetic control of immune response to, 1204
Nucleic acid, ribo-, tumor virus, endogenous, autogenous immunity to, identification of antibody reactivity to select viral antigens, 1568
— acid, ribo-, viruses, mammalian C-type, common cell surface antigen associated with, 925
Nucleotide levels, cyclic, enzyme release, and phagocytosis, effects of autonomic agents on, hormonal control of lysosomal enzyme release from human neutrophils, 1395
Nude mice grafted with co-isogenic or allogeneic thymuses, activity of

host-derived T cells which differentiate in, 1215

OCTAPEPTIDE-HEME coupled to Fab, preparation and characterization of immunoelectron microscope tracer consisting of, 208

Ontogeny of B lymphocytes, 1125, 1142

Organs, lymphoid, peripheral, identification of novel cell type in, functional properties in vitro, 380

—, lymphoid, peripheral, identification of novel cell type in, functional properties in vivo, 1431

Oxidized and reduced ascorbate, uptake and reduction by human leukocytes, 1084

PANENCEPHALITIS, sclerosing subacute, partial characterization of blocking factor in plasma and spinal fluid of patients with, 902

Parabiosed mice, immunological tolerance, dissociation between in vivo and in vitro reactivity in, 767

Peyer's patch lymphoid cells, induction of humoral antibody and cell-mediated allograft reactions, 398

— patch lymphoid cells, lipopolysaccharide is thymus dependent, 407

Phagocytes, mononuclear, in vitro synthesis and secretion of lysozome by, 1228

Phagocytic stimulus, segmental response of macrophage plasma membrane to, 323

Phagocytosis, enzyme release, and cyclic nucleotide levels, effects of autonomic agents on, hormonal control of lysosomal enzyme release from human neutrophils, 1395

Phosphorylcholine, genetics of new IgV_H (T15 idiotype) marker in mouse regulating natural antibody to, 983

Phosphorylcholine-binding receptors on lymphoid cells, specificity, class, and idiotype of, clonal nature of immune response to, 58

Phytomitogens and allogeneic lymphocytes, cellular interactions in proliferative response of human T and B lymphocytes to, 1553

Plasma membrane, macrophage, segmental response to phagocytic stimulus, 323

— and spinal fluid of patients with subacute sclerosing panencephalitis, partial characterization of blocking factor in, 902

Plasminogen activator, secretion by stimulated macrophages, 834

— activators formed by neoplastic human cell cultures, properties, 1317

Pneumococcal polysaccharide, Type III, in mice, genetic control of antibody response to, relationship between IgM immunoglobulin levels and ability to give IgM antibody response, 1499

Polymorphism, new, in mice, electrophoretic mobility and agglutinability of red blood cells, 313

Polypeptide, collagen-like synthetic, thymus independence of, need for thymus and bone marrow cell cooperation in immune response to gelatin, 148

Polysaccharide, pneumococcal, Type III, in mice, genetic control of antibody response to, relationship between IgM immunoglobulin levels and ability to give IgM antibody response, 1499

Polysaccharides, streptococcal group, in mice, genetics of restricted antibodies to, strain differences of isoelectric focusing spectra of group A hyperimmune antisera, 1513

Pressure, blood, increased, and mortality in spontaneously hypertensive rat, 617

Priming, antigen, recent, effect on adoptive immune responses, 1

Properdin factor D, characterization of active site and isolation of precursor form, 355
— role in alternate pathway of complement activation, 44
Protein, amyloid fibril, new, third class, 773
— conjugates, hapten autologous, lymphocyte proliferation in vitro induced by, 732
—, human, acute phase, found in association with rubella virus infection, characterization, 497
Proteins, myeloma, immunochemical analysis of idiotypes with specificity for levan or dextran, 137
—, myeloma, reactive with dextrans or with fructosan and on human antilevans, immunochemical studies on, 159
Pyrogens, leukocytic, human, two, demonstration and characterization, 1369

RAJI cells, lymphoblastoid, human, receptor for soluble C3 and C3b on, properties and biological significance, 696
Receptor antibody, anti-T-cell, inhibition of specific cell-mediated cytotoxicity by, 888
—, antigen, removal and regeneration after binding of antigen to lymphocytes, 1110
—, complement, lymphocytes, *H-2* linkage of gene controlling rate of appearance of, 1142
— for soluble C3 and C3b on human lymphoblastoid, Raji, cells, properties and biological significance, 696
Receptor-carrying cells, characterization by use of antibodies to alloantibodies, 877
Receptors, C′3 and Fc, effect of B-cell mitogens on lymphocyte subpopulations possessing, 969
—, complement, and surface immunoglobulin, relative rates of appearance of lymphocytes bearing, 1125
— for aggregated IgG on thymocytes, thymus-derived and bone marrow-derived lymphocytes, 1175
Receptors, lymphocyte, responding in vitro to hapten autologous protein conjugates, specificity and nature of, 732
—, phosphorylcholine-binding, on lymphoid cells, specificity, class, and idiotype of, clonal nature of immune response, 58
—, specific, general method for isolation and recovery of B cells bearing, 761
Reed-Sternberg and other cells in Hodgkin's disease, immunohistochemical demonstration of IgG in, 1077
Renal. *See* Kidney.
Resistance, cell-mediated, to *Listeria monocytogenes*, effect of sea star coelomocyte extract on, 820
—, genetic, to Friend virus leukemia in mice, mechanisms, 1093
Ribonucleic acid tumor virus, endogenous, autogenous immunity to, identification of antibody reactivity to select viral antigens, 1568
— acid viruses, mammalian C-type, common cell surface antigen associated with, 925
Rubella virus infection, characterization of human acute phase protein found in association with, 497

SALMONELLA infection, acute, cytokinetics of monocytosis in, 264
Sarcoma virus, Moloney, evidence of suppressor cell activity in mice spleens bearing primary tumors induced by, 1473
Sclerosing panencephalitis, subacute, partial characterization of blocking factor in plasma and spinal fluid of patients with, 902
Sea star coelomocyte extract, effect on cell-mediated resistance to *Listeria monocytogenes*, 820
Secretion of plasminogen activator by stimulated macrophages, 834
—, synthesis, and cell surface expression of immunoglobulin in mu-

rine thoracic duct lymphocytes, 1013
Secretion and synthesis, in vitro, of lysozyme by mononuclear phagocytes, 1228
—, synthesis, and shedding of immunoglobulin by lymphoid cells of germ-free mice, 862
Secretory component, free and bound, localization in human intestinal epithelial cells, model for assembly of secretory IgA, 629
Sensitized lymphocytes, specifically, feedback inhibition of, 543
Serum factor in chronic hypocomplementemic nephritis distinct from immunoglobulins and activating alternate complement pathway, 1249
Sex-linked defect in cross-linking of collagen and elastin associated with mottled locus in mice, 180
Sodium chloride, toxicity, increased blood pressure and mortality in spontaneously hypertensive rat, 617
Soluble antigens, defined, cytotoxic immune cells with specificity for, 93
— C3 and C3b on human lymphoblastoid, Raji, cells, receptor for, properties and biological significance, 696
Specific binding of IgE, evidence in interaction of IgE with basophilic leukemia cells, 600
— cell-mediated cytotoxicity, inhibition by anti-T-cell receptor antibody, 888
— cytotoxicity, antibody as mediator of, 93
—, hapten-, inactivated, precursors of antibody-forming cells, effects of allogeneic cell interactions and enzymatic digestion with trypsin on, 1446
— heterologous enhancement of immune responses, 439
— immune responses, mitogenic activation of B cells results in, 74
— receptors, general method for isolation and recovery of B cells bearing, 761
Specific T-cell helper function, regulatory influence of bacteria lipopolysaccharide on, 24
—, tumor-, cytotoxicity to urinary bladder carcinoma, lymphoid cells mediating, separation of effector population using surface marker, 457
— unresponsiveness of circulating lymphocytes from mice primed with heterologous erythrocytes, 1
Specifically and cross-immunized hamsters, cell-mediated immunity against *Besnoitia* and *Toxoplasma* in, and in cultures, 560
— sensitized lymphocytes, feedback inhibition of, 543
Specificities, cross-idiotypic, similarities in light chains of anti-γ-globulins showing, 128
—, *H-2D*^{d}, genetic analysis of responses to, in mixed lymphocyte culture responses of mice, 851
Specificity, class, and idiotype of phosphorylcholine-binding receptors on lymphoid cells, clonal nature of immune response, 58
— for defined soluble antigens, cytotoxic immune cells with, 93
— for levan or dextran, immunochemical analysis of idiotypes of myeloma proteins with, 137
— and nature of lymphocyte receptors responding in vitro to hapten autologous protein conjugates, 732
Spinal fluid and plasma of patients with subacute sclerosing panencephalitis, partial characterization of blocking factor in, 902
Spleens, mice, bearing primary tumors induced by Moloney sarcoma virus, evidence of suppressor cell activity in, 1473
Splenic lymphocytes from tumor-bearing mice, mixed lymphocyte reactivity against normal cells by, 224, 230
Splenocytes, murine, new method for study of synthesis, intracellular transport, and exteriorization of cell surface immunoglobulin in, 1599

Staphylococcal nuclease, genetic control of immune response to, 1204
Stem cells, hemopoietic, and myeloid progenitors, antigenic differences between, 1621
— cells, semiallogeneic, transplantation into immunodeficient chicks, complete restoration of bursa-dependent immune system after, 1344
Stimulation, cellular, lectin-induced, interaction of cholera toxin and toxin derivatives with lymphocytes, binding properties and interference with, 801
—, immuno-, humoral, increased uptake of [^{125}I]iododeoxyuridine and [^{3}H]thymidine into TNP cells treated with anti-TNP antibody, 367
— of antibody production to hapten 2,4-dinitrobenzene by affinity labeled murine lymphoid cells, 943
— of cell motility by anti-Ig and lack of relationship to capping, 295
Streptococcal, group A, cell wall substance, changes in cellular enzyme levels and extracellular release of lysosomal acid hydrolases in macrophages exposed to, 1262
— group polysaccharides in mice, genetics of restricted antibodies to, strain differences of isoelectric focusing spectra of group A hyperimmune antisera, 1513
^{89}Strontium-sensitive effector cells responsible for rejection of bone marrow allografts, role of, 1093
Synergy among responding lymphoid cells in one-way mixed lymphocyte reaction, interaction between two types of thymus-dependent cells, 1488
Syngeneic and semisyngeneic systems, autoimmune-like activity in, 230

TARGET antigens, cellular distribution in alloantiserum-induced inhibition of immune response gene product function, 661
— antigens, genetic analysis in alloantiserum-induced inhibition of immune response gene product function, 679
Thoracic duct lymphocytes, murine, synthesis, secretion, and cell surface expression of immunoglobulin in, 1013
Thymic suppressor cell, separation of subpopulations with suppressor activity, 13
Thymidine, ^{3}H-labeled, and ^{125}I-labeled iododeoxyuridine, increased uptake into TNP cells treated with anti-TNP antibody, 367
Thymocytes, thymus-derived and bone marrow-derived lymphocytes, presence of receptors for aggregated IgG on, 1175
Thymoma cell immunoglobulin, inhibition of T-cell-dependent immune response in vitro by, 785
Thymus cells, absence of MLR-stimulatory, M-locus product from, 1628
Thymus, dual regulatory role in maturation of immune response, 108
— humoral factor, hormone-like activity on induction of immune competence in lymphoid cells, 193
— independence of collagen-like synthetic polypeptide and of collagen, need for thymus and bone marrow cell cooperation in immune response to gelatin, 148
— lipid and brain inhibition of anti-brain-associated θ-cytotoxicity, 445
Thymus-dependent antigen, induction of immunological tolerance to in absence of thymus-derived cells, 1303
— cells, two types, interaction between, synergy among responding lymphoid cells in one-way mixed lymphocyte reaction, 1488
— lipopolysaccharide, Peyer's patch lymphoid cells, 407
Thymus-derived, anti-, cell receptor antibody, inhibition of specific cell-mediated cytotoxicity by, 888
— and bone marrow-derived lymphocytes, activation in vitro, 24

Thymus-derived and bone marrow-derived lymphocytes, human, cellular interactions in proliferative response to phytomitogens and allogeneic lymphocytes, 1553
— and bone marrow-derived lymphocytes, roles in termination of unresponsiveness to autologous thyroglobulin, 643
— and bone marrow-derived lymphocytes, and thymocytes, presence of receptors for aggregated IgG on, 1175
— cell activities in mixed lymphocyte reaction and two graft-vs.-host assays, dissociation of, 721
— cell lines, human malignant, Epstein-Barr virus-negative, 1070
— cell-dependent immune response, inhibition in vitro by thymoma cell immunoglobulin, 785
— cell-mediated immunity, potentiation by selective suppression of antibody formation with cyclophosphamide, 1529
— cells, host-derived, which differentiate in nude mice grafted with co-isogenic or allogeneic thymuses, activity of, 1215
— cells, induction of immunological tolerance to thymus-dependent antigen in absence of, 1303
— cells, influence of dose and route of antigen injection on immunological induction of, 528
— cells, modifying effect of BCG on immunological induction of, 1540
— lymphocytes, activated, binding of aggregated γ-globulin to, 1002
— lymphocytes, two populations immune to lymphocytic choriomeningitis virus, 1329
— and marrow-derived lymphocytes, origin of cell surface immunoglobulin of, 479
Thymus-independent immunocyte triggering, mitogenic activation of B cells results in specific immune responses, 74
Thymuses, co-isogenic or allogeneic, activity of host-derived T cells which differentiate in nude mice grafted with, 1215
Thyroglobulin, autologous, roles of T and B lymphocytes in termination of unresponsiveness to, 643
Tolerance, immunologic, partial, induction in rats and progressive loss of cellular antigenicity in Gross virus lymphoma, 1382
—, immunological, dissociation between in vivo and in vitro reactivity in parabiosed mice, 767
—, immunological, in bone marrow-derived lymphocytes, 1446, 1464
—, immunological, to thymus-dependent antigen, induction in absence of thymus-derived cells, 1303
Toxicity of sodium chloride, increased blood pressure and mortality in spontaneously hypertensive rat, 617
Toxin, cholera, and toxin derivatives, interaction with lymphocytes, binding properties and interference with lectin-induced cellular stimulation, 801
Toxoplasma and *Besnoitia* in specifically and cross-immunized hamsters and in cultures, cell-mediated immunity against, 560
— *gondii*, effect of normal and activated human macrophages on, 1154
Tracer, immunoelectron microscope, consisting of heme-octapeptide coupled to Fab, preparation and characterization, 208
—study, ultrastructural, of altered functional properties of renal glomerulus in autologous immune complex nephritis, 1283
Transplantation of semiallogeneic stem cells into immunodeficient chicks, complete restoration of bursa-dependent immune system after, 1344
—, renal, in diabetes mellitus in rats, 793
Triggering, immunocyte, thymus-independent, mitogenic activation of

B cells results in specific immune responses, 74
2,4,6-Trinitrophenyl cells treated with anti-TNP antibody, increased uptake of [^{125}I]iododeoxyuridine and [^{3}H]thymidine into, 367
Trypsin, enzymatic digestion with and allogeneic cell interactions, effects on inactivated hapten-specific precursors of antibody-forming cells, 1446
Tumor cells, parental strain, vigorous immune responsiveness induced in F_1 mice by, 224
— virus, RNA, endogenous, autogenous immunity to, identification of antibody reactivity to select viral antigens, 1568
Tumor-bearing mice, mixed lymphocyte reactivity against normal cells by splenic lymphocytes from, 224, 230
Tumor-specific cytotoxicity to urinary bladder carcinoma, lymphoid cells mediating, separation of effector population using surface marker, 457
Tumors, primary, induced by Moloney sarcoma virus, evidence of suppressor cell activity in mice spleens bearing, 1473

URINARY bladder carcinoma, lymphoid cells mediating tumor-specific cytotoxicity to, separation of effector population using surface marker, 457

VENOM factor, cobra, and other mitogens, complement-dependent B-cell activation by, 337
Viral antigens, identification of antibody reactivity to, autogenous immunity to endogenous RNA tumor virus, 1568
— gene products, C-type, of virus-negative mouse cells, isolation and characterization, 427
Virus, choriomeningitis, lymphocytic, two populations of T lymphocytes immune to, 1329
—, encephalomyocarditis, genetic influences affecting occurrence of diabetes mellitus-like disease in mice infected with, 414
— infection, rubella, characterization of human acute phase protein found in association with, 497
—, leukemia, Friend, in mice, mechanisms of genetic resistance to, 1093
—, lymphoma, Gross, induction of partial immunologic tolerance in rats and progressive loss of cellular antigenicity in, 1382
—, sarcoma, Moloney, evidence of suppressor cell activity in mice spleens bearing primary tumors induced by, 1473
—, tumor, RNA, endogenous, autogenous immunity to, identification of antibody reactivity to select viral antigens, 1568
Virus-negative, Epstein-Barr, human malignant T-cell lines, 1070
— mouse cells, isolation and characterization of C-type viral gene products of, 427
Viruses, RNA, mammalian C-type, common cell surface antigen associated with, 925